

Moldy Hay for Horses¹

Haymaking conditions this spring have been poor. Much hay has been rained on or left lying in the field for prolonged time periods due to cool and humid conditions

which reduced drying rates. The long drying periods with high humidity allowed field growth of mold on the hay.

Poor drying weather has also meant that some hay was put up wetter than usual and mold growth occurred in storage. With wet weather and high humidity, normal drying in storage may not occur and hay can retain elevated levels of moisture allowing mold growth. Mold will grow on hay without preservative added at moisture levels above 14% to 15%. The mold growth produces heat and can result in large amounts of dry matter and TDN (total digestible nutrient) loss – a loss of carbohydrates and binding of proteins. In some cases, heating can be great enough to cause spontaneous combustion and fire. Drying of stored hay (moisture loss) is enhanced by ventilation, creation of air spaces between bales, allowing ample head space above a stack of bales in a barn for moisture to evaporate since moisture tends to move up and out the top of a stack of bales, avoidance of other wet products in the same area (if enclosed storage), reduced size of stacks, alternated direction of stacking, and not placing tarp directly over a stack in the field because that traps moisture.



Molds commonly found in hay include *Alternaria*, *Aspergillus*, *Cladosporium*, *Fusarium*, *Mucor*, *Penicillium*, and *Rhizopus*. These molds **can produce spores that cause respiratory problems**, especially in horses and, **under some conditions, will produce mycotoxins**.

Horses are particularly sensitive to dust from mold spores and can get a respiratory disease similar to asthma in humans called Recurrent Airway Obstruction (RAO), commonly referred to as heaves. A horse with RAO will have a normal temperature and a good appetite, but will often have decreased exercise tolerance, excessive tearing of the eyes, coughing and nasal discharge. The horse will appear winded. Some horses are highly allergic to certain mold spores while others seem to be unaffected. Even among horses with symptoms, there seems to be a variation to their sensitivity level. To decrease exposure, have the horse spend more time outside at pasture than inside and feed the horse outside if possible. Some other ways to reduce dust exposure are as follows:

- Use dust-free bedding such as shredded paper or rubber mats.
- Don't feed dusty and moldy hay and grains.
- Keep your horse out of the stable when you are cleaning to reduce exposure to dust.
- Feed hay outside to minimize dust problems.
- Soak dusty hay for 30 minutes before feeding so that the horse can eat it while it's wet.
- Store hay away from your horse as much as possible and ensure any hay in the vicinity is kept dry to reduce mold.
- If the horse is housed indoors, ensure that there is good, draught-free ventilation through the stable.

¹ Written by Dan Undersander, University of Wisconsin (djunders@wisc.edu); Marvin Hall, The Pennsylvania State University (mhh2@psu.edu); Richard Leep, Michigan State University (leep@msu.edu); Glenn Shewmaker, University of Idaho (gshew@uidaho.edu); Don Westerhaus, Kemin AgriFoods North America (Don.Westerhaus@kemin.com), and Lon Whitlow, North Carolina State University (lon_whitlow@ncsu.edu).

Sometimes mold spores are counted on moldy feeds to obtain an indication of the extent of molding and relative risks in feeding them. **Table 1** contains classification of risks at various mold spore counts.

While most molds do not produce mycotoxins, the presence of mold indicates the possibility of mycotoxin presence and animals being fed moldy hay should be watched carefully for mycotoxin symptoms.

Table 1. Feeding Risks^a at Various Mold Spore Counts	
Mold Spore count per gram	Feeding Risk and Cautions
Under 500,000	Relatively low Risk
½ to 1 million	Relatively Safe
1 to 2 million	Feed with Caution
2 to 3 million	Closely observe animals and performance
3 to 5 million	Dilute with other feeds
Over 5 million	Discontinue feeding
^a Risks refer primarily to effect of mold per se without regard to possible mycotoxin content. Data from Richard S. Adams, Kenneth B. Kephart, Virginia A. Ishler, Lawrence J. Hutchinson, and Gregory W. Roth. 1993. Mold and mycotoxin problems in livestock feeding. The Pennsylvania State University.	

Mycotoxins effects on animals:

- 1) intake reduction or feed refusal;
- 2) reduced nutrient absorption and impaired metabolism, including altered digestion and microbial growth, diarrhea, intestinal irritation, reduced production, lower fertility, abortions, lethargy, and increased morbidity;
- 3) alterations in the endocrine and exocrine systems;
- 4) suppression of the immune system which predisposes livestock to many diseases and may increase milk somatic cell count. A suppressed immune system may also cause lack of response to medications and failure of vaccine programs;
- 5) cellular death causing organ damage.

If you have mold in hay, watch for the symptoms mentioned above. If hay is dusty (from mold spores) take care in feeding to sensitive animals and those in areas with poor ventilation. If mycotoxin symptoms are observed, check with a nutritionist to make sure the ration is properly balanced and possibly with a veterinarian to eliminate other disease/health problems. Quick test kits (ELISA kits) are available (listing at <http://www.ces.ncsu.edu/gaston/Agriculture/mycotoxins/mycotest.html>) to determine presence of a limited number mycotoxins but they can give false positives. Some forage testing laboratories will provide other mycotoxin tests. Often, the best strategy is to remove a suspected mycotoxin-contaminated feedstuff from the diet and see if symptoms disappear. If mycotoxins are present, the feedstuff can often be fed at a diluted rate and/or with approved feed additives.

In summary:

- ◆ Most molds are harmless producing spores and dust but not producing known mycotoxins.
- ◆ Many of the commonly diagnosed mycotoxins^a are produced in the field prior to harvest.
- ◆ If a mycotoxin problem is suspected, a comprehensive review of herd nutrition and health is essential - e.g. herd problems blamed on mycotoxins may be other disorders or nutritional issues. Diagnosing a mycotoxin problem is difficult and often involves the elimination of other possible factors.
- ◆ The physical dust problem associated with moldy forage can be reduced by feeding in a well ventilated area, mixing with a high moisture feed or wetting the hay, but these will not reduce mycotoxins if present.